

STANLEY SYMMETRIC FUNCTIONS AND QUIVER VARIETIES

ANDERS SKOVSTED BUCH

1. INTRODUCTION

The purpose of this paper is to show a connection between Stanley symmetric functions and the formula for quiver varieties given in [5]. Recall that a simple reflection in the symmetric group S_m is a transposition that interchanges two consecutive integers. A *reduced word* for a permutation $w \in S_m$ is a tuple of simple reflections $(\tau_1, \tau_2, \dots, \tau_\ell)$ with $\ell = \ell(w)$ the length of w , such that $w = \tau_1 \tau_2 \cdots \tau_\ell$. Stanley asked how many reduced words does a permutation w have.

To answer this question, Stanley [17] defined a power series $F_w(x)$ in infinitely many variables x_1, x_2, \dots ; it is homogeneous of degree $\ell = \ell(w)$, has non-negative integer coefficients, and the number of reduced words for w is the coefficient in $F_w(x)$ of the monomial $x_1 x_2 \cdots x_\ell$. Stanley then proved that this power series is symmetric. This implies that it can be written in the basis of Schur functions:

$$(1) \quad F_w(x) = \sum_{\lambda \vdash \ell} \alpha_{w\lambda} s_\lambda(x)$$

where the sum is over all partitions λ of ℓ and the coefficients $\alpha_{w\lambda}$ are integers. Since the coefficient of $x_1 x_2 \cdots x_\ell$ in a Schur function $s_\lambda(x)$ is equal to the number f^λ of standard Young tableaux of shape λ (see e.g. [15] or [10]), it follows that the number of reduced words for w is given as

$$\sum_{\lambda \vdash \ell} \alpha_{w\lambda} f^\lambda.$$

The constants f^λ are considered well understood, so only a description of the coefficients $\alpha_{w\lambda}$ remained to be found. Stanley credits Edelman and Greene for proving that these coefficients are non-negative [6] (see also [14]). Fomin and Greene have shown that $\alpha_{w\lambda}$ is equal to the number of semistandard Young tableaux T of shape λ , such that the column word of T is a reduced word for w [7]. Another useful fact is that $\alpha_{w^{-1}\lambda} = \alpha_{w\lambda'}$ where λ' is the conjugate of λ [14], [16, (7.22)].

Stanley's symmetric function is known to be a limit of Schubert polynomials $\mathfrak{S}_w(x)$ defined by Lascoux and Schützenberger [13], [16]. For $n \in \mathbb{N}$, let $1^n \times w \in S_{n+m}$ denote the shifted permutation which acts as the identity on $1, \dots, n$ and maps i to $w(i - n) + n$ for $n + 1 \leq i \leq n + m$. If one specializes to finitely many variables x_1, x_2, \dots, x_N , then

$$(2) \quad F_w(x_1, \dots, x_N, 0, 0, \dots) = \mathfrak{S}_{1^n \times w^{-1}}(x_1, \dots, x_N, 0, 0, \dots)$$

for all sufficiently large n [16, (7.18)].

Date: February 8, 2008.

The formula for quiver varieties given in [5] specializes to a formula for the double Schubert polynomial $\mathfrak{S}_w(x; y)$ for the permutation $w \in S_m$:

$$(3) \quad \mathfrak{S}_w(x; y) = \sum c_w(a, b, \lambda) y_2^{a_2} \cdots y_{m-1}^{a_{m-1}} (-x_2)^{b_2} \cdots (-x_{m-1})^{b_{m-1}} s_\lambda(x/y).$$

The sum is over exponents a_2, \dots, a_{m-1} and b_2, \dots, b_{m-1} and a partition λ . The coefficients $c_w(a, b, \lambda)$ are special cases of a large class of generalized Littlewood-Richardson coefficients, which are conjectured to be non-negative and given by a generalized Littlewood-Richardson rule [5].

The main result in this paper is that the coefficient $c_w(0, 0, \lambda)$ (corresponding to zero exponents) is equal to Stanley's coefficient $\alpha_{w^{-1}\lambda}$. In this way, (3) writes a Schubert polynomial $\mathfrak{S}_w(x)$ as a symmetric polynomial equal to Stanley's symmetric function for w^{-1} plus a non-symmetric polynomial.

In Section 2 and Section 3 we review the results of [5]. In Section 4 we prove the identity $\alpha_{w^{-1}\lambda} = c_w(0, 0, \lambda)$ and use this to give a new proof of Stanley's result [17] that the symmetric function $F_{w_0}(x)$ for the longest permutation w_0 in S_m is equal to the Schur function $s_\lambda(x)$ for the staircase partition $\lambda = (m-1, m-2, \dots, 1)$. In Section 5 we use geometry of degeneracy loci to prove a generalization of the well known formula for a Schubert polynomial of a product of two permutations. Finally, in Section 6, we discuss the relations to the conjectured generalized Littlewood-Richardson rule in [5].

We thank S. Fomin, W. Fulton, and F. Sottile for helpful discussions.

2. A FORMULA FOR QUIVER VARIETIES

Let X be a non-singular complex variety and $E_0 \rightarrow E_1 \rightarrow E_2 \rightarrow \cdots \rightarrow E_n$ a sequence of vector bundles and vector bundle maps over X . A set of *rank conditions* for this sequence is a collection $r = (r_{ij})$ of non-negative integers, for $0 \leq i < j \leq n$. Let $\Omega_r(E_\bullet) \subset X$ be the locus where each composite map $E_i \rightarrow E_j$ has rank at most r_{ij} :

$$\Omega_r(E_\bullet) = \{x \in X \mid \text{rank}(E_i(x) \rightarrow E_j(x)) \leq r_{ij} \ \forall i < j\}.$$

This locus determines a cohomology class $[\Omega_r(E_\bullet)]$ in the cohomology ring $H^*(X) = H^*(X; \mathbb{Z})$ of X .

For convenience we set $r_{ii} = \text{rank}(E_i)$. We will require that the rank conditions $r = (r_{ij})$ can *occur*, i.e. there should exist a sequence of vector spaces and linear maps $V_0 \rightarrow V_1 \rightarrow \cdots \rightarrow V_n$ such that $\dim(V_i) = r_{ii}$ and $\text{rank}(V_i \rightarrow V_j) = r_{ij}$ for all $i < j$. This is equivalent to demanding $r_{ij} \leq \min(r_{i,j-1}, r_{i+1,j})$ for $i < j$ and $r_{ij} - r_{i,j-1} - r_{i+1,j} + r_{i+1,j-1} \geq 0$ for $j - i \geq 2$ [1].

If the locus $\Omega_r(E_\bullet)$ is not empty, its codimension is at most

$$d(r) = \sum_{i < j} (r_{i,j-1} - r_{ij})(r_{i+1,j} - r_{ij}).$$

Furthermore, this codimension is obtained for generic choices of bundle maps $E_i \rightarrow E_{i+1}$ [5], [12], [2]. The main result of [5] gives a formula for the cohomology class of the degeneracy locus $\Omega_r(E_\bullet)$, when it has this expected codimension $d(r)$.

To explain this formula, we need some notation. Let $\Lambda = \mathbb{Z}[h_1, h_2, \dots]$ be the ring of symmetric functions. The variable h_i can be identified with the complete symmetric function of degree i in variables x_i , $i \in \mathbb{N}$. If $I = (a_1, a_2, \dots, a_p)$ is a

sequence of integers, define the Schur function $s_I \in \Lambda$ to be the determinant of the $p \times p$ matrix whose $(i, j)^{\text{th}}$ entry is h_{a_i+j-i} :

$$s_I = \det(h_{a_i+j-i})_{1 \leq i, j \leq p}.$$

(Here one sets $h_0 = 1$ and $h_{-q} = 0$ for $q > 0$.) A Schur function is always equal to either zero, or plus or minus a Schur function for a partition λ :

$$s_I = \begin{cases} 0 \\ \pm s_\lambda. \end{cases}$$

This follows from interchanging the rows of the matrix defining s_I . Furthermore, the Schur functions given by partitions form a basis for the ring of symmetric functions [15], [10].

If E is a vector bundle over X , let $c_i(E) \in H^{2i}(X)$ denote its i^{th} Chern class. Given two vector bundles E and F of ranks e and f over X one can define a ring homomorphism

$$\Lambda \rightarrow H^*(X)$$

which maps h_i to the coefficient of t^i in the formal power series expansion of the quotient

$$\frac{c_t(E^\vee)}{c_t(F^\vee)} = \frac{1 - c_1(E)t + \cdots + (-1)^e c_e(E)t^e}{1 - c_1(F)t + \cdots + (-1)^f c_f(F)t^f}.$$

We let $s_\lambda(F - E) \in H^*(X)$ denote the image of s_λ by this map. (If E and F have Chern roots x_1, \dots, x_e and y_1, \dots, y_f respectively, the notation $s_\lambda(y/x) = s_\lambda(F - E)$ is also common.)

When the degeneracy locus $\Omega_r(E_\bullet)$ has its expected codimension $d(r)$, its cohomology class is equal to a linear combination of products of Schur polynomials in differences of consecutive bundles:

$$[\Omega_r(E_\bullet)] = \sum_{\mu} c_\mu(r) s_{\mu_1}(E_1 - E_0) s_{\mu_2}(E_2 - E_1) \cdots s_{\mu_n}(E_n - E_{n-1}).$$

Here the sum is over sequences of partitions $\mu = (\mu_1, \mu_2, \dots, \mu_n)$. The coefficients $c_\mu(r)$ are integer constants depending on the rank conditions and the sequence μ . They are determined by a combinatorial algorithm which we will describe next.

Start by arranging the rank conditions $r = (r_{ij})$ in a *rank diagram*:

$$\begin{array}{ccccccc} E_0 & \rightarrow & E_1 & \rightarrow & E_2 & \rightarrow & \cdots & \rightarrow & E_n \\ r_{00} & & r_{11} & & r_{22} & & \cdots & & r_{nn} \\ & r_{01} & & r_{12} & & \cdots & & & r_{n-1,n} \\ & & r_{02} & & \cdots & & & & r_{n-2,n} \\ & & & \ddots & & & & & \\ & & & & r_{0n} & & & & \end{array}$$

In this diagram, replace each small triangle of numbers

$$\begin{array}{cc} r_{i,j-1} & r_{i+1,j} \\ & r_{ij} \end{array}$$

by a rectangle R_{ij} with $r_{i+1,j} - r_{ij}$ rows and $r_{i,j-1} - r_{ij}$ columns.

$$R_{ij} = \begin{array}{c} \begin{array}{|c|c|c|c|} \hline & & & \\ \hline & & & \\ \hline & & & \\ \hline & & & \\ \hline \end{array} \\ r_{i,j-1} - r_{ij} \end{array} \begin{array}{c} r_{i+1,j} - r_{ij} \\ \\ \\ \end{array}$$

These rectangles are arranged in a *rectangle diagram*:

$$\begin{array}{ccccccc} R_{01} & & R_{12} & & \cdots & & R_{n-1,n} \\ & R_{02} & & \cdots & & R_{n-2,n} & \\ & & \ddots & & & & \\ & & & R_{0n} & & & \end{array}$$

The information carried by the rank conditions is very well represented in this diagram. First, the expected codimension $d(r)$ for the locus $\Omega_r(E_\bullet)$ is equal to the total number of boxes in the rectangle diagram. Furthermore, the condition that the rank conditions can occur is equivalent to saying that the rectangles get narrower when one travels south-west, while they get shorter when one travels south-east. Finally, the algorithm that computes the coefficients $c_\mu(r)$ depends only on the rectangle diagram.

We will define this algorithm by constructing an element P_r in the n^{th} tensor power of the ring of symmetric functions $\Lambda^{\otimes n}$, so that

$$P_r = \sum_{\mu} c_\mu(r) s_{\mu_1} \otimes \cdots \otimes s_{\mu_n}.$$

This is done by induction on n . When $n = 1$ (corresponding to a sequence of two vector bundles), the rectangle diagram has only one rectangle $R = R_{01}$. In this case we set

$$P_r = s_R \in \Lambda^{\otimes 1},$$

where R is identified with the partition for which it is the Young diagram. This case recovers the Giambelli-Thom-Porteous formula.

If $n \geq 2$, we let \bar{r} denote the bottom n rows of the rank diagram. Then \bar{r} is a valid set of rank conditions, so by induction we can assume that

$$(4) \quad P_{\bar{r}} = \sum_{\mu} c_\mu(\bar{r}) s_{\mu_1} \otimes \cdots \otimes s_{\mu_{n-1}}$$

is a well defined element of $\Lambda^{\otimes n-1}$. Now P_r is obtained from $P_{\bar{r}}$ by replacing each basis element $s_{\mu_1} \otimes \cdots \otimes s_{\mu_{n-1}}$ in (4) with the sum

$$\sum_{\substack{\sigma_1, \dots, \sigma_{n-1} \\ \tau_1, \dots, \tau_{n-1}}} \left(\prod_{i=1}^{n-1} c_{\sigma_i \tau_i}^{\mu_i} \right) s_{\begin{array}{|c|} \hline R_{01} \\ \hline \end{array} \begin{array}{|c|} \hline \sigma_1 \\ \hline \end{array}} \otimes \cdots \otimes s_{\begin{array}{|c|} \hline R_{i-1,i} \\ \hline \end{array} \begin{array}{|c|} \hline \sigma_i \\ \hline \end{array}} \otimes \cdots \otimes s_{\begin{array}{|c|} \hline R_{n-1,n} \\ \hline \end{array} \begin{array}{|c|} \hline \tau_{n-1} \\ \hline \end{array}}.$$

This sum is over all partitions $\sigma_1, \dots, \sigma_{n-1}$ and $\tau_1, \dots, \tau_{n-1}$ such that σ_i has fewer rows than $R_{i-1,i}$ and each Littlewood-Richardson coefficient $c_{\sigma_i \tau_i}^{\mu_i}$ is non-zero. A diagram consisting of a rectangle $R_{i-1,i}$ with (the Young diagram of) a partition σ_i attached to its right side, and τ_{i-1} attached beneath should be interpreted as the sequence of integers giving the number of boxes in each row of this diagram.

It can happen that the rectangle $R_{i-1,i}$ is empty, since the number of rows or columns can be zero. If the number of rows is zero, then σ_i is required to be empty, and the diagram is the Young diagram of τ_{i-1} . If the number of columns is zero, then the algorithm requires that the length of σ_i is at most equal to the number of rows $r_{ii} - r_{i-1,i}$, and the diagram consists of σ_i in the top $r_{ii} - r_{i-1,i}$ rows and τ_{i-1} below this, possibly with some zero-length rows in between.

3. SCHUBERT POLYNOMIALS

Let $w \in S_{m+1}$ be a permutation, and let E_\bullet be a sequence of bundles over X

$$F_1 \subset F_2 \subset \cdots \subset F_m \rightarrow G_m \twoheadrightarrow G_{m-1} \twoheadrightarrow \cdots \twoheadrightarrow G_1$$

consisting of a full flag with a general map to a dual full flag. Define the locus

$$\Omega_w = \{x \in X \mid \text{rank}(F_q(x) \rightarrow G_p(x)) \leq r_w(p, q) \ \forall p, q\}$$

where $r_w(p, q) = \#\{i \leq p \mid w(i) \leq q\}$. Fulton has proved [8] that the cohomology class of this locus is given by the double Schubert polynomial defined by Lascoux and Schützenberger [13]:

$$[\Omega_w] = \mathfrak{S}_w(x_1, \dots, x_m; y_1, \dots, y_m)$$

where $x_i = c_1(\ker(G_i \rightarrow G_{i-1}))$ and $y_i = c_1(F_i/F_{i-1})$. Now $\Omega_w = \Omega_r(E_\bullet)$ where $r = (r_{ij})$ are the obvious rank conditions. This means that the double Schubert polynomial becomes a special case of the quiver formula:

$$\begin{aligned} \mathfrak{S}_w(x; y) &= [\Omega_r(E_\bullet)] \\ &= \sum c_\mu(r) s_{\mu_1}(F_2 - F_1) \cdots s_{\mu_{m-1}}(F_m - F_{m-1}) \cdot s_{\mu_m}(G_m - F_m) \cdot \\ &\quad s_{\mu_{m+1}}(G_{m-1} - G_m) \cdots s_{\mu_{2m-1}}(G_1 - G_2) \end{aligned}$$

As noted in [5], significant simplifications can be made by using the equalities

$$s_\lambda(F_{i+1} - F_i) = s_\lambda(y_{i+1}) = \begin{cases} y_{i+1}^a & \text{if } \lambda = (a) \text{ is a row with } a \text{ boxes} \\ 0 & \text{otherwise} \end{cases}$$

and

$$s_\lambda(G_i - G_{i+1}) = s_\lambda(0/x_{i+1}) = \begin{cases} (-x_{i+1})^b & \text{if } \lambda = (1^b) \text{ is a column with } b \text{ boxes} \\ 0 & \text{otherwise.} \end{cases}$$

Using this and the fact that $s_\lambda(G_m - F_m)$ is the super-symmetric Schur polynomial $s_\lambda(x/y)$ in the variables x_1, \dots, x_m and y_1, \dots, y_m , we obtain a formula

$$(5) \quad \mathfrak{S}_w(x; y) = \sum c_w(a, b, \lambda) y_2^{a_2} \cdots y_m^{a_m} (-x_2)^{b_2} \cdots (-x_m)^{b_m} s_\lambda(x/y).$$


The sum is over exponents a_2, \dots, a_m and b_2, \dots, b_m , and a single partition λ , and $c_w(a, b, \lambda)$ is the coefficient $c_\mu(r)$ for the sequence of partitions

$$\mu = ((a_2), \dots, (a_m), \lambda, (1^{b_m}), \dots, (1^{b_2})).$$

Example 1. For the permutation $w = 2431$ we get the rank diagram

$$\begin{array}{cccccccc} F_1 & \subset & F_2 & \subset & F_3 & \rightarrow & G_3 & \twoheadrightarrow & G_2 & \twoheadrightarrow & G_1 \\ 1 & & 2 & & 3 & & 3 & & 2 & & 1 \\ & & 1 & & 2 & & 2 & & 2 & & 1 \\ & & & & 1 & & 1 & & 1 & & \\ & & & & 0 & & 1 & & 1 & & \\ & & & & & & 0 & & 1 & & \\ & & & & & & & & 0 & & \end{array}$$

This means that the rectangle diagram for $1 \times w$ is obtained by adding a rim of empty rectangles to the sides of the rectangle diagram for w .



Similarly, one obtains the rectangle diagram for $1^n \times w$ by adding n rims of empty rectangles to the rectangle diagram for w .

Let $P_r \in \Lambda^{\otimes 2m-1}$ be the element associated to the rank conditions $r = (r_{ij})$ for w . The above comparison of rectangle diagrams then shows that $1^n \times w$ corresponds to the element

$$\underbrace{1 \otimes \cdots \otimes 1}_n \otimes P_r \otimes \underbrace{1 \otimes \cdots \otimes 1}_n \in \Lambda^{\otimes 2m+2n-1}.$$

By (5) this gives us

$$\mathfrak{S}_{1^n \times w}(x; y) = \sum c_w(a, b, \lambda) y_{2+n}^{a_2} \cdots y_{m+n}^{a_m} (-x_{2+n})^{b_2} \cdots (-x_{m+n})^{b_m} s_\lambda(x/y)$$

where $s_\lambda(x/y)$ is in variables x_1, \dots, x_{m+n} and y_1, \dots, y_{m+n} .

Now restrict to two fixed sets of variables x_1, \dots, x_N and y_1, \dots, y_M , setting $x_i = y_j = 0$ for $i > N$ and $j > M$. When $n \geq \max(N-1, M-1)$, the only non-zero terms in the above expression for $\mathfrak{S}_{1^n \times w}$ are those with all exponents a_i and b_i equal to zero. Since this Schubert polynomial is homogeneous of degree equal to the length of w , the partitions λ occurring in these terms all have weight $\ell(w)$. This proves:

Theorem. *Let $w \in S_{m+1}$ and fix two sets of variables x_1, \dots, x_N and y_1, \dots, y_M . When $n \geq \max(N-1, M-1)$, the double Schubert polynomial $\mathfrak{S}_{1^n \times w}$ in these variables is given by*

$$\mathfrak{S}_{1^n \times w}(x_1, \dots, x_N, 0, \dots, 0; y_1, \dots, y_M, 0, \dots, 0) = \sum_{\lambda \vdash \ell(w)} c_w(0, 0, \lambda) s_\lambda(x/y).$$

Comparing with equations (1) and (2) we obtain (since $\alpha_{w\lambda} = \alpha_{w^{-1}\lambda'}$):


Corollary. *Stanley's coefficient $\alpha_{w\lambda}$ is equal to $c_w(0, 0, \lambda')$.*

Thus the formula (3) writes a Schubert polynomial as a symmetric part equal to Stanley's symmetric function plus additional non-symmetric terms. For example, if $w = 2431$ as in the above example, we have $F_w(x) = s_{\square\square}(x)$.

The identity $\alpha_{w\lambda'} = \alpha_{w^{-1}\lambda}$ becomes a special case of the identity $c_{w^{-1}}(b, a, \lambda') = c_w(a, b, \lambda)$, which in turn follows from the formula $c_{\mu^\vee}(r^\vee) = c_\mu(r)$ of [5]. Here r^\vee are the rank conditions obtained by mirroring the rank diagram for $r = (r_{ij})$ in a vertical line, so $r_{ij}^\vee = r_{n-j, n-i}$, and μ^\vee is the sequence (μ'_n, \dots, μ'_1) of conjugate partitions in the opposite order.

Example 2. Let $w_0 = m \cdots 21$ be the longest permutation in S_m . Then we have $r_{w_0}(p, q) = \max(p+q-m, 0)$. The rectangle diagram associated to w_0 therefore

has exactly i non-empty rectangles in the i^{th} row for $1 \leq i \leq m-1$, and these are centered around the middle. All other rectangles are empty.



We will use this diagram to compute Stanley's symmetric function for w_0 . The idea is that in order for the algorithm to produce a contribution to F_{w_0} , all boxes must travel north-west until they meet the last non-empty rectangle in this direction, and from that point they must travel north-east.

Let $P_r^{(k)}$ denote the element in $\Lambda^{\otimes 2m-3-k}$ given by this rectangle diagram with the top k rows removed. In particular $P_r^{(0)} = P_r$ is the element associated to the whole diagram. The terms s_λ in Stanley's symmetric function F_{w_0} are in 1-1 correspondence with terms in P_r of the form

$$\underbrace{1 \otimes \cdots \otimes 1}_{m-2} \otimes s_\lambda \otimes \underbrace{1 \otimes \cdots \otimes 1}_{m-2}.$$

One may check that, in order for a term

$$s_{\mu_1} \otimes \cdots \otimes s_{\mu_{2m-3-k}}$$

in $P_r^{(k)}$ to contribute to F_{w_0} , the partition μ_i must be empty if the i^{th} rectangle in the $k+1^{\text{st}}$ row of the rectangle diagram is empty, while it must have length at most one unless the i^{th} rectangle is the leftmost non-empty rectangle in the $k+1^{\text{st}}$ row. To be precise, the term $s_{\mu_1} \otimes \cdots \otimes s_{\mu_{2m-3-k}}$ contributes to F_{w_0} only if μ_i has length at most one for $i \neq m-1-k$, and is empty when $i \leq m-2-k$ and when $i \geq m$. The reason is that all rectangles in the diagram have height at most one, which means that any σ_i in the algorithm can have length at most one. So if any μ_i has two or more rows, boxes are forced to the right, creating a new partition with too many rows if $i \neq m-1-k$.

An examination of the algorithm then shows that $P_r^{(k)}$ contains only one such term, with coefficient 1. This term has μ_{m-1-k} equal to the staircase partition with $m-1-k$ rows, $\mu_{m-1-k} = (m-1-k, \dots, 2, 1)$, while any other non-empty partitions μ_i is a single row with $m-1-k$ boxes, $\mu_i = (m-1-k)$.

Taking $k=0$, we see that $F_{w_0} = s_{(m-1, m-2, \dots, 2, 1)}$. This was first proved by Stanley [17], and implies that the number of reduced words for w_0 is equal to the number of standard tableaux on the staircase partition with $m-1$ rows.

5. REDUNDANT RANK CONDITIONS AND PRODUCTS OF PERMUTATIONS

Suppose we are given a sequence of bundles $E_0 \rightarrow E_1 \rightarrow \cdots \rightarrow E_n$ and a set of rank conditions $r = \{r_{ij}\}$ for this sequence. The degeneracy locus $\Omega_r(E_\bullet)$ is then the subset of points $x \in X$ over which the maps on fibers satisfy all of the inequalities


$$\text{rank}(E_i(x) \rightarrow E_j(x)) \leq r_{ij}$$

for $i < j$. Some of these inequalities may be redundant in the sense that they follow from other inequalities. It is easy to see that the inequality involving the number

r_{ij} is redundant if and only if this number is equal to one of $r_{i,j-1}$ or $r_{i+1,j}$. In other words, the inequality involving r_{ij} is necessary if and only if the rectangle R_{ij} is not empty.

Now suppose there are integers $0 \leq p \leq q \leq n$ such that the rectangle R_{ij} is empty whenever exactly one of i and j is in the interval $[p, q]$. In this case the degeneracy locus $\Omega_r(E_\bullet)$ is the (scheme-theoretic) intersection of two larger loci $\Omega_{r'}(E'_\bullet)$ and $\Omega_{r''}(E''_\bullet)$ for the sequences $E'_\bullet : E_p \rightarrow E_{p+1} \rightarrow \cdots \rightarrow E_q$ and $E''_\bullet : E_1 \rightarrow \cdots \rightarrow E_{p-1} \rightarrow E_{q+1} \rightarrow \cdots \rightarrow E_n$, where r' and r'' are the restrictions of the rank conditions $r = \{r_{ij}\}$ to these sequences. We will say that E'_\bullet is an independent subsequence. Note that if $p = q$, the bundle E_p is redundant and can be removed from the sequence E_\bullet without changing $\Omega_r(E_\bullet)$. This special case was described in [5].

When E'_\bullet is an independent subsequence, the rectangle diagram for the rank conditions r' simply consists of the rectangles R_{ij} for $p \leq i < j \leq q$, while the rectangle diagram for r'' contains the remaining non-empty rectangles.



This in particular means that the expected codimensions of $\Omega_{r'}(E'_\bullet)$ and $\Omega_{r''}(E''_\bullet)$ add up to that of $\Omega_r(E_\bullet)$. If all of these loci have their expected codimensions, then we get the equality $[\Omega_r(E_\bullet)] = [\Omega_{r'}(E'_\bullet)] \cdot [\Omega_{r''}(E''_\bullet)]$ in the cohomology ring of X . To see this, note at first that both $\Omega_{r'}(E'_\bullet)$ and $\Omega_{r''}(E''_\bullet)$ are Cohen-Macaulay [12] (see also [11, Lemma A.2]). If $f : \Omega_{r'}(E'_\bullet) \hookrightarrow X$ is the inclusion, we therefore get

$$[\Omega_r(E_\bullet)] = f_*[f^{-1}(\Omega_{r''}(E''_\bullet))] = f_*f^*[\Omega_{r''}(E''_\bullet)] = [\Omega_{r'}(E'_\bullet)] \cdot [\Omega_{r''}(E''_\bullet)].$$

This means that the formula P_r satisfies


$$(6) \quad P_r = \underbrace{(1 \otimes \cdots \otimes 1)}_p \otimes P_{r'} \otimes \underbrace{(1 \otimes \cdots \otimes 1)}_{n-q} \cdot \Phi_p^{q-p+2}(P_{r''})$$

where multiplication is performed factor-wise, and Φ_p^k denotes the k -fold coproduct expansion of the p^{th} factor of its arguments, i.e.

$$\begin{aligned} \Phi_p^k(s_{\mu_1} \otimes \cdots \otimes s_{\mu_p} \otimes \cdots \otimes s_{\mu_\ell}) = \\ \sum_{\sigma_1, \dots, \sigma_k} c_{\sigma_1, \dots, \sigma_k}^{\mu_p} s_{\mu_1} \otimes \cdots \otimes s_{\mu_{p-1}} \otimes s_{\sigma_1} \otimes \cdots \otimes s_{\sigma_k} \otimes s_{\mu_{p+1}} \otimes \cdots \otimes s_{\mu_\ell}. \end{aligned}$$

We will apply this to study the Schubert polynomial of a product of two permutations. If $w \in S_m$ and $u \in S_n$ are permutations, define the product $w \times u \in S_{m+n}$ to be the permutation which maps i to $w(i)$ if $1 \leq i \leq m$, while $m+i$ is mapped

to $m + u(i)$ for $1 \leq i \leq n$. The rank diagram for this permutation is equal to that of $1^m \times u$, except the bottom $2m - 2$ rows are replaced by the rank diagram for w . The diamond of non-empty rectangles in the rectangle diagram for $w \times u$ is therefore split into a top part containing the diamond of rectangles for u and a bottom part with the diamond of rectangles for w .



Given a sequence of bundles consisting of a full flag of length $m + n - 1$ followed by a full dual flag of the same length as in Section 3, we deduce that the locus $\Omega_{w \times u}$ is the intersection of the loci Ω_w and $\Omega_{1^m \times u}$. We therefore recover the well known formula [16, (4.6)]

$$(7) \quad \mathfrak{S}_{w \times u} = \mathfrak{S}_w \cdot \mathfrak{S}_{1^m \times u}$$

for the Schubert polynomial of a product of two permutations. This immediately implies Stanley's identity $F_{w \times u} = F_w \cdot F_u$ [17]. Note that the same argument shows that (7) also holds for Fulton's universal Schubert polynomials [9].

6. RELATIONS TO A CONJECTURED LITTLEWOOD-RICHARDSON RULE

In this final section we will discuss relations with Stanley symmetric functions of a generalized Littlewood-Richardson rule which is conjectured in [5]. We will need the notions of (semistandard) Young tableaux and multiplication of tableaux, see for example [10].

A *tableau diagram* for a set of rank conditions $r = (r_{ij})$ is a filling of all boxes in the corresponding rectangle diagram with integers, so that each rectangle R_{ij} becomes a tableau T_{ij} . Furthermore, it is required that the entries of each tableau T_{ij} are strictly larger than the entries in tableaux above T_{ij} in the diagram, within 45 degree angles. These are the tableaux T_{kl} with $i \leq k < l \leq j$ and $(k, l) \neq (i, j)$.

A *factor sequence* for a tableau diagram with n rows is a sequence of tableaux (W_1, \dots, W_n) , which is obtained as follows: If $n = 1$ then the only factor sequence is the sequence (T_{01}) containing the only tableau in the diagram. When $n \geq 2$, a factor sequence is obtained by first constructing a factor sequence (U_1, \dots, U_{n-1}) for the bottom $n - 1$ rows of the tableau diagram, and choosing arbitrary factorizations of the tableaux in this sequence:

$$U_i = P_i \cdot Q_i.$$

Then the sequence


$$(W_1, \dots, W_n) = (T_{01} \cdot P_1, Q_1 \cdot T_{12} \cdot P_2, \dots, Q_{n-1} \cdot T_{n-1,n})$$

is a factor sequence for the whole tableau diagram.

In [5] it is conjectured that the coefficient $c_\mu(r)$ is equal to the number of different factor sequences (W_1, \dots, W_n) for any fixed tableau diagram for the rank conditions r , such that W_i has shape μ_i for each i . This conjecture has been proved (using an involution of Fomin) when all the rectangles in the fourth row of the rectangle diagram and below are empty, and no two non-empty rectangles in the third row are neighbors [4].

If $r = (r_{ij})$ are the rank conditions given by a permutation w , then the conjecture implies that Stanley's coefficient $\alpha_{w\lambda}$ is equal to the number of different tableaux W of shape λ' , for which $(\emptyset, \dots, \emptyset, W, \emptyset, \dots, \emptyset)$ is a factor sequence. Thus a proof of the general conjecture will give a new proof that Stanley's coefficients are non-negative, as well as an interesting way to compute them.

Example 3. Let $w = 2143 \dots (2p)(2p-1) \in S_{2p}$ for some $p > 0$. Then the rectangle diagram for w has a 1×1 rectangle in the middle of row $4i+1$ for $0 \leq i \leq p-1$. All other rectangles are empty.



A tableau diagram is obtained by filling the numbers $1, 2, \dots, p$ in these boxes. It is easy to see that a sequence $(\emptyset, \dots, \emptyset, W, \emptyset, \dots, \emptyset)$ is a factor sequence for this diagram if and only if W is a standard tableau with p boxes. Therefore the conjecture predicts that Stanley's symmetric function is given by

$$F_w = \sum_{\lambda \vdash p} f^\lambda s_\lambda.$$

This can be confirmed using Stanley's formula $F_{w \times u} = F_w \cdot F_u$ [17]. Let $\sigma = 21 \in S_2$. Then $w = \sigma \times \dots \times \sigma$ (p times), which implies that

$$F_w = (F_\sigma)^p = (s_{\square})^p = \sum_{\lambda \vdash p} f^\lambda s_\lambda.$$

We thank F. Sottile for showing us a different proof of this fact.

Using a criterion for factor sequences given in [4], one may also prove that the conjectured Littlewood-Richardson rule gives the correct prediction for Stanley's symmetric function of a longest permutation w_0 .

In general, Stanley's symmetric function F_w is known to have a minimal term $s_{\lambda(w)}$ and a maximal term $s_{\mu(w)}$, both occurring with coefficient one. If $w \in S_{m+1}$, define

$$r_p(w) = \#\{q \mid q < p \text{ and } w(q) > w(p)\}$$

for $1 \leq p \leq m+1$, and let $\lambda(w)$ be the partition obtained by arranging the numbers $r_1(w), \dots, r_{m+1}(w)$ in decreasing order. Let $\mu(w)$ be the conjugate of the partition $\lambda(w^{-1})$. Then $\alpha_{w, \lambda(w)} = \alpha_{w, \mu(w)} = 1$, and any partition λ with $\alpha_{w\lambda} \neq 0$ is between $\lambda(w)$ and $\mu(w)$ in the dominance order [17].

Let $\{T_{ij}\}_{1 \leq i < j \leq 2m}$ be a tableau diagram for (the rank conditions given by) w . There are two extremal ways to form a factor sequence $(\emptyset, \dots, \emptyset, W, \emptyset, \dots, \emptyset)$ for this diagram. The first is to make all factorizations of inductive factor sequences (U_1, \dots, U_k) be “rightward” whenever possible. This means that when factoring U_i into $U_i = P_i \cdot Q_i$, we take $P_i = \emptyset$ and $Q_i = U_i$ for $i \neq m$ while we take $P_m = U_m$ and $Q_m = \emptyset$ (if $k \geq m$). The middle tableau in the final factor sequence then is

$$W_{\text{right}} = T_m \cdot T_{m+1} \cdot \dots \cdot T_{2m-1}$$

where

$$T_j = T_{0j} \cdot T_{1j} \cdot \dots \cdot T_{m-1,j}.$$

Note that each tableau T_j has only one column. If we set $p = 2m + 1 - j$ and $q = w^{-1}(i + 2)$ then T_{ij} is non-empty if and only if $q < p$ and $w(q) > w(p)$. It follows that T_j has exactly $r_p(w)$ boxes.

We claim that W_{right} has shape $\lambda(w)'$, corresponding to the maximal term of $F_{w^{-1}}$. It is enough to show that if T_l and T_j both have a box in row t and $l < j$, then the box in T_l is smaller than the one in T_j . To prove this, let the t^{th} box in T_l come from T_{kl} and the t^{th} box in T_j come from T_{ij} . If the box in T_{kl} is not smaller than the box in T_{ij} then $k < i$. Now since the tableau T_{il} must be as wide as T_{kl} and as tall as T_{ij} , this tableau T_{il} can't be empty. Similarly, if T_{hj} corresponds to a box over T_{ij} in T_j , then T_{hl} gives a corresponding box in T_l . This shows that the boxes corresponding to T_{kl} and T_{ij} in T_l and T_j was not in the same row, a contradiction.

Similarly one can show that the tableau obtained by “leftward” factorizations,

$$W_{\text{left}} = (T_{0,m} \cdot T_{0,m+1} \cdot \dots \cdot T_{0,2m-1}) \cdot (T_{1,m} \cdot \dots \cdot T_{1,2m-1}) \cdot \dots \cdot (T_{m-1,m} \cdot \dots \cdot T_{m-1,2m-1}),$$

has shape $\mu(w)'$.

REFERENCES

- [1] S. Abeasis and A. Del Fra, *Degenerations for the representations of an equioriented quiver of type A_m* , Boll. Un. Mat. Ital. Suppl. (1980), no. 2, 157–171.
- [2] S. Abeasis, A. Del Fra, and H. Kraft, *The geometry of representations of A_m* , Math. Ann. **256** (1981), 401–418.
- [3] A. S. Buch, *Combinatorics of Degeneracy Loci*, A dissertation submitted to the faculty of the Division of the Physical Sciences, The University of Chicago, 1999.
- [4] ———, *On a conjectured formula for quiver varieties*, preprint, 1999.
- [5] A. S. Buch and W. Fulton, *Chern class formulas for quiver varieties*, Invent. math. **135** (1999), 665–687.
- [6] P. Edelman and C. Greene, *Balanced tableaux*, Adv. in Math. **63** (1987), 42–99.
- [7] S. Fomin and C. Greene, *Noncommutative Schur functions and their applications*, Discrete Math. **193** (1998), no. 1-3, 179–200.
- [8] W. Fulton, *Flags, Schubert polynomials, degeneracy loci, and determinantal formulas*, Duke Math. J. **65** (1992), 381–420.
- [9] ———, *Universal Schubert polynomials*, Duke Math. J. **96** (1999), no. 3, 575–594.
- [10] ———, *Young tableaux*, London Mathematical Society Student Texts, vol. 35, Cambridge University Press, 1997.
- [11] W. Fulton and P. Pragacz, *Schubert varieties and degeneracy loci*, Springer-Verlag, Berlin, 1998.
- [12] V. Lakshmibai and P. Magyar, *Degeneracy schemes and Schubert varieties*, preprint, 1997.
- [13] A. Lascoux and M.-P. Schützenberger, *Polynômes de Schubert*, C. R. Acad. Sci. Paris Sér. I Math. **294** (1982), 447–450.
- [14] ———, *Structure de Hopf de l’anneau de cohomologie et de l’anneau de Grothendieck d’une variété de drapeaux*, C. R. Acad. Sci. Paris Sér. I Math. **295** (1982), 629–633.

- [15] I. G. Macdonald, *Symmetric functions and Hall polynomials*, Oxford University Press, 1979, 1995.
- [16] ———, *Notes on Schubert polynomials*, Laboratoire de Combinatoire et d'Informatique Mathématique, Université du Québec à Montréal, 1991.
- [17] R. P. Stanley, *On the number of reduced decompositions of elements of Coxeter groups*, European J. Combin. **5** (1984), 359–372.

MASSACHUSETTS INSTITUTE OF TECHNOLOGY, BUILDING 2, ROOM 248, 77 MASSACHUSETTS AVENUE, CAMBRIDGE, MA 02139

E-mail address: `abuch@math.mit.edu`